L Number	Hits	Search Text	DB	Time stamp
2	66	(delta adj1 (fos or FosB)) or FosB	USPAT;	2003/05/06 08:13
			US-PGPUB;	
			EPO;	}
	_		DERWENT	
1	1	(delta adj1 (fos or FosB))	USPAT;	2003/05/06 08:13
			US-PGPUB;	
		•	EPO;	
_			DERWENT	
3	66	FosB	USPAT;	2003/05/06 08:14
			US-PGPUB;	
			EPO;	
4	24	FosB and (skeletal or bone or osteo?)	DERWENT	2003/05/06 00:15
"	24	rosb and (skeretar of bone of osteo?)	USPAT;	2003/05/06 08:15
			US-PGPUB; EPO;	
-			DERWENT	
5	4	FosB.clm.	USPAT;	2003/05/06 08:15
1	•	1 00D 1 01	US-PGPUB;	2003/03/00 08:13
			EPO;	
			DERWENT	

From:

Woitach, Joseph

Sent:

Tuesday, May 06, 2003 10:55 AM

To: Subject: STIC-ILL references

Hello.

Could you send me the following references:

Cell 1991 Feb 22;64(4):751-9 Nakabeppu Y, Nathans D.

A naturally occurring truncated form of FosB that inhibits Fos/Jun transcriptional activity.

Mol Cell Biol 1993 Jul;13(7):4157-66

Nakabeppu Y, Oda S, Sekiguchi M.

Proliferative activation of quiescent Rat-1A cells by delta FosB

J Neurochem 1997 Jan;68(1):272-9

Bing G, Wang W, Qi Q, Feng Z, Hudson P, Jin L, Zhang W, Bing R, Hong JS.

Long-term expression of Fos-related antigen and transient expression of delta FosB associated with seizures in the

rat hippocampus and striatum.

Eur J Neurosci 1998 May;10(5):1743-51

Porcella A, Gessa GL, Pani L.

Delta9-tetrahydrocannabinol increases sequence-specific AP-1 DNA-binding activity and Fosrelated antigens in

the rat brain.

Brain Res Mol Brain Res 1998 Jan;53(1-2):41-52

Perez-Otano I. Mandelzys A. Morgan Jl.

MPTP-Parkinsonism is accompanied by persistent expression of a delta-FosB-like protein in dopaminergic

pathways.

J Neurosci. 1997 Jul 15;17(14):5407-15

Mandelzys A. Gruda MA, Bravo R, Morgan Jl.

Absence of a persistently elevated 37 kDa fos-related antigen and AP-1-like DNA-binding activity in the brains of kainic acid-treated fosB null

mice.

Brain Res Mol Brain Res 1997 Jul;47(1-2):31-43

Hollen KM, Nakabeppu Y, Davies SW.

Changes in expression of delta FosB and the Fos family proteins following NMDA receptor activation in the rat

striatum.

Oncogene 1995 Apr 6;10(7):1343-51

Oda S, Nishida J, Nakabeppu Y, Sekiguchi M.

Stabilization of cyclin E and cdk2 mRNAs at G1/S transition in Rat-1A cells emerging from the G0 state.

Neuron 1994 Nov;13(5):1235-44

Hope BT, Nye HE, Kelz MB, Self DW, ladarola MJ, Nakabeppu Y, Duman RS, Nestler EJ. Induction of a long-lasting AP-1 complex composed of altered Fos-like proteins in brain by chronic cocaine and

other chronic treatments.

WP351.565 (NPL) 444235

A CONTRACTOR OF THE PROPERTY O

THE PERSON NAMED IN COLUMN TWO IS NOT THE PERSON NAMED IN THE PE

Woitach, Joseph

Tuesday, May 06, 2003 10:55 AM

To:

STIC-ILL

Subject:

From:

Sent:

references

Hello.

Could you send me the following references:

Cell 1991 Feb 22;64(4):751-9 Nakabeppu Y, Nathans D.

A naturally occurring truncated form of FosB that inhibits Fos/Jun transcriptional activity.

Mol Cell Biol 1993 Jul;13(7):4157-66

Nakabeppu Y, Oda S, Sekiguchi M.

Proliferative activation of quiescent Rat-1A cells by delta FosB

J Neurochem 1997 Jan;68(1):272-9

Bing G, Wang W, Qi Q, Feng Z, Hudson P, Jin L, Zhang W, Bing R, Hong JS.

Long-term expression of Fos-related antigen and transient expression of delta FosB associated with seizures in the

rat hippocampus and striatum.

Eur J Neurosci 1998 May:10(5):1743-51

Porcella A. Gessa GL. Pani L.

Delta9-tetrahydrocannabinol increases sequence-specific AP-1 DNA-binding activity and Fosrelated antigens in

the rat brain.

Brain Res Mol Brain Res 1998 Jan:53(1-2):41-52

Perez-Otano I, Mandelzys A, Morgan JI.

MPTP-Parkinsonism is accompanied by persistent expression of a delta-FosB-like protein in dopaminergic

pathways.

J Neurosci. 1997 Jul 15;17(14):5407-15

Mandelzys A, Gruda MA, Bravo R, Morgan Jl.

Absence of a persistently elevated 37 kDa fos-related antigen and AP-1-like DNA-binding activity in the brains of kainic acid-treated fosB null

mice.

Brain Res Mol Brain Res 1997 Jul;47(1-2):31-43

Hollen KM, Nakabeppu Y, Davies SW.

Changes in expression of delta FosB and the Fos family proteins following NMDA receptor activation in the rat

striatum.

Oncogene 1995 Apr 6;10(7):1343-51

Oda S, Nishida J, Nakabeppu Y, Sekiguchi M.

Stabilization of cyclin E and cdk2 mRNAs at G1/S transition in Rat-1A cells emerging from the G0 state.

Neuron 1994 Nov;13(5):1235-44

Hope BT, Nye HE, Kelz MB, Self DW, ladarola MJ, Nakabeppu Y, Duman RS, Nestler EJ. Induction of a long-lasting AP-1 complex composed of altered Fos-like proteins in brain by chronic cocaine and

other chronic treatments.

AP 356-2-B23 Adonis

From: Sent: Woitach, Joseph

Sent: To: Tuesday, May 06, 2003 10:55 AM

Subject:

STIC-ILL references

Hello.

Could you send me the following references:

Cell 1991 Feb 22;64(4):751-9 Nakabeppu Y, Nathans D.

A naturally occurring truncated form of FosB that inhibits Fos/Jun transcriptional activity.

Mol Cell Biol 1993 Jul;13(7):4157-66

Nakabeppu Y, Oda S, Sekiguchi M.

Proliferative activation of quiescent Rat-1A cells by delta FosB

J Neurochem 1997 Jan;68(1):272-9

Bing G, Wang W, Qi Q, Feng Z, Hudson P, Jin L, Zhang W, Bing R, Hong JS.

Long-term expression of Fos-related antigen and transient expression of delta FosB associated with seizures in the

rat hippocampus and striatum.

Eur J Neurosci 1998 May;10(5):1743-51

Porcella A, Gessa GL, Pani L.

Delta9-tetrahydrocannabinol increases sequence-specific AP-1 DNA-binding activity and Fosrelated antigens in

the rat brain.

Brain Res Mol Brain Res 1998 Jan;53(1-2):41-52

Perez-Otano I, Mandelzys A, Morgan Jl.

MPTP-Parkinsonism is accompanied by persistent expression of a delta-FosB-like protein in dopaminergic

pathways.

J Neurosci. 1997 Jul 15;17(14):5407-15

Mandelzys A, Gruda MA, Bravo R, Morgan Jl.

Absence of a persistently elevated 37 kDa fos-related antigen and AP-1-like DNA-binding activity in the brains of kainic acid-treated fosB null

_mice.

Brain Res Mol Brain Res 1997 Jul;47(1-2):31-43

Hollen KM, Nakabeppu Y, Davies SW.

Changes in expression of delta FosB and the Fos family proteins following NMDA receptor activation in the rat

striatum.

Oncogene 1995 Apr 6:10(7):1343-51

Oda S, Nishida J, Nakabeppu Y, Sekiguchi M.

Stabilization of cyclin E and cdk2 mRNAs at G1/S transition in Rat-1A cells emerging from the G0 state.

Neuron 1994 Nov;13(5):1235-44

Hope BT, Nye HE, Kelz MB, Self DW, ladarola MJ, Nakabeppu Y, Duman RS, Nestler EJ. Induction of a long-lasting AP-1 complex composed of altered Fos-like proteins in brain by chronic cocaine and

other chronic treatments.

356. 2.B73

The second secon

1,

From:

Woitach, Joseph

Sent:

Tuesday, May 06, 2003 10:55 AM

To: Subject: STIC-ILL references

Hello,

Could you send me the following references:

Cell 1991 Feb 22;64(4):751-9 Nakabeppu Y, Nathans D.

A naturally occurring truncated form of FosB that inhibits Fos/Jun transcriptional activity.

Mol Cell Biol 1993 Jul; 13(7): 4157-66

Nakabeppu Y, Oda S, Sekiguchi M.

Proliferative activation of quiescent Rat-1A cells by delta FosB

J Neurochem 1997 Jan;68(1):272-9

Bing G, Wang W, Qi Q, Feng Z, Hudson P, Jin L, Zhang W, Bing R, Hong JS.

Long-term expression of Fos-related antigen and transient expression of delta FosB associated with seizures in the

rat hippocampus and striatum.

Eur J Neurosci 1998 May;10(5):1743-51

Porcella A. Gessa GL. Pani L.

Delta9-tetrahydrocannabinol increases sequence-specific AP-1 DNA-binding activity and Fosrelated antigens in

the rat brain.

Brain Res Mol Brain Res 1998 Jan;53(1-2):41-52

Perez-Otano I, Mandelzys A, Morgan Jl.

MPTP-Parkinsonism is accompanied by persistent expression of a delta-FosB-like protein in dopaminergic

pathways.

J Neurosci. 1997 Jul 15;17(14):5407-15

Mandelzys A, Gruda MA, Bravo R, Morgan Jl.

Absence of a persistently elevated 37 kDa fos-related antigen and AP-1-like DNA-binding activity in the brains of kainic acid-treated fosB null mice.

Brain Res Mol Brain Res 1997 Jul;47(1-2):31-43 Hollen KM, Nakabeppu Y, Davies SW.

Changes in expression of delta FosB and the Fos family proteins following NMDA receptor activation in the rat

striatum.

Oncogene 1995 Apr 6:10(7):1343-51

Oda S, Nishida J, Nakabeppu Y, Sekiguchi M.

Stabilization of cyclin E and cdk2 mRNAs at G1/S transition in Rat-1A cells emerging from the G0 state.

Neuron 1994 Nov;13(5):1235-44

Hope BT, Nye HE, Kelz MB, Self DW, ladarola MJ, Nakabeppu Y, Duman RS, Nestler EJ. Induction of a long-lasting AP-1 complex composed of altered Fos-like proteins in brain by chronic cocaine and

other chronic treatments.

Woitach, Joseph

Tuesday, May 06, 2003 10:55 AM

Sent: To:

From:

STIC-ILL

Subject:

references

Hello,

Could you send me the following references:

Cell 1991 Feb 22;64(4):751-9 Nakabeppu Y, Nathans D.

A naturally occurring truncated form of FosB that inhibits Fos/Jun transcriptional activity.

NOS

944 496

Mol Cell Biol 1993 Jul;13(7):4157-66

Nakabeppu Y, Oda S, Sekiguchi M.

Proliferative activation of guiescent Rat-1A cells by delta FosB

J Neurochem 1997 Jan:68(1):272-9

Bing G, Wang W, Qi Q, Feng Z, Hudson P, Jin L, Zhang W, Bing R, Hong JS.

Long-term expression of Fos-related antigen and transient expression of delta FosB associated with seizures in the

rat hippocampus and striatum.

Eur J Neurosci 1998 May:10(5):1743-51

Porcella A, Gessa GL, Pani L.

Delta9-tetrahydrocannabinol increases sequence-specific AP-1 DNA-binding activity and Fosrelated antigens in

the rat brain.

Brain Res Mol Brain Res 1998 Jan;53(1-2):41-52

Perez-Otano I. Mandelzys A. Morgan Jl.

MPTP-Parkinsonism is accompanied by persistent expression of a delta-FosB-like protein in dopaminergic

pathways.

J Neurosci. 1997 Jul 15:17(14):5407-15

Mandelzys A, Gruda MA, Bravo R, Morgan JI.

Absence of a persistently elevated 37 kDa fos-related antigen and AP-1-like DNA-binding activity in the brains of kainic acid-treated fosB null mice.

Brain Res Mol Brain Res 1997 Jul:47(1-2):31-43

Hollen KM, Nakabeppu Y, Davies SW.

Changes in expression of delta FosB and the Fos family proteins following NMDA receptor activation in the rat

striatum.

Oncogene 1995 Apr 6:10(7):1343-51

Oda S, Nishida J, Nakabeppu Y, Sekiguchi M.

Stabilization of cyclin E and cdk2 mRNAs at G1/S transition in Rat-1A cells emerging from the G0 state.

Neuron 1994 Nov:13(5):1235-44

Hope BT, Nye HE, Kelz MB, Self DW, ladarola MJ, Nakabeppu Y, Duman RS, Nestler EJ. Induction of a long-lasting AP-1 complex composed of altered Fos-like proteins in brain by chronic cocaine and

other chronic treatments.

QP35-1.557

THE RESERVE TO SHARE THE PARTY OF THE PARTY

TO THE RESIDENCE OF THE PROPERTY OF THE PROPER

The second of th

From:

Woitach, Joseph

Sent:

Tuesday, May 06, 2003 10:55 AM

To: Subject: STIC-ILL references

Hello,

Could you send me the following references:

Cell 1991 Feb 22;64(4):751-9 Nakabeppu Y, Nathans D.

A naturally occurring truncated form of FosB that inhibits Fos/Jun transcriptional activity.

Mol Cell Biol 1993 Jul;13(7):4157-66

Nakabeppu Y, Oda S, Sekiguchi M.

Proliferative activation of quiescent Rat-1A cells by delta FosB

J Neurochem 1997 Jan;68(1):272-9

Bing G, Wang W, Qi Q, Feng Z, Hudson P, Jin L, Zhang W, Bing R, Hong JS.

Long-term expression of Fos-related antigen and transient expression of delta FosB

rat hippocampus and striatum.

Eur J Neurosci 1998 May;10(5):1743-51

Porcella A, Gessa GL, Pani L.

Delta9-tetrahydrocannabinol increases sequence-specific AP-1 DNA-binding activity and Fosrelated antigens in

the rat brain.

Brain Res Mol Brain Res 1998 Jan;53(1-2):41-52

Perez-Otano I, Mandelzys A, Morgan Jl.

MPTP-Parkinsonism is accompanied by persistent expression of a delta-FosB-like protein in dopaminergic

pathways.

J Neurosci. 1997 Jul 15;17(14):5407-15

Mandelzys A, Gruda MA, Bravo R, Morgan Jl.

Absence of a persistently elevated 37 kDa fos-related antigen and AP-1-like DNA-binding activity in the brains of kainic acid-treated fosB null mice.

Brain Res Mol Brain Res 1997 Jul;47(1-2):31-43

Hollen KM, Nakabeppu Y, Davies SW.

Changes in expression of delta FosB and the Fos family proteins following NMDA receptor activation in the rat

striatum.

Oncogene 1995 Apr 6:10(7):1343-51

Oda S, Nishida J, Nakabeppu Y, Sekiguchi M.

Stabilization of cyclin E and cdk2 mRNAs at G1/S transition in Rat-1A cells emerging from the G0 state.

Neuron 1994 Nov;13(5):1235-44

Hope BT, Nye HE, Kelz MB, Self DW, ladarola MJ, Nakabeppu Y, Duman RS, Nestler EJ. Induction of a long-lasting AP-1 complex composed of altered Fos-like proteins in brain by chronic cocaine and

other chronic treatments.

Woitach, Joseph

From: Sent:

To:

Tuesday, May 06, 2003 10:55 AM

Subject:

STIC-ILL references

Hello,

Could you send me the following references:

Cell 1991 Feb 22;64(4):751-9 Nakabeppu Y, Nathans D.

A naturally occurring truncated form of FosB that inhibits Fos/Jun transcriptional activity.

DH506.M6 Mi

The state of the s

The second secon

Mol Cell Biol 1993 Jul; 13(7): 4157-66

Nakabeppu Y, Oda S, Sekiguchi M.

Proliferative activation of quiescent Rat-1A cells by delta FosB

J Neurochem 1997 Jan;68(1):272-9

Bing G, Wang W, Qi Q, Feng Z, Hudson P, Jin L, Zhang W, Bing R, Hong JS.

Long-term expression of Fos-related antigen and transient expression of delta FosB associated with seizures in the

rat hippocampus and striatum.

Eur J Neurosci 1998 May:10(5):1743-51

Porcella A. Gessa GL. Pani L.

Delta9-tetrahydrocannabinol increases sequence-specific AP-1 DNA-binding activity and Fosrelated antigens in

the rat brain.

Brain Res Mol Brain Res 1998 Jan; 53(1-2):41-52

Perez-Otano I, Mandelzys A, Morgan JI.

MPTP-Parkinsonism is accompanied by persistent expression of a delta-FosB-like protein in dopaminergic

pathways.

J Neurosci. 1997 Jul 15:17(14):5407-15

Mandelzys A, Gruda MA, Bravo R, Morgan Jl.

Absence of a persistently elevated 37 kDa fos-related antigen and AP-1-like DNA-binding activity in the brains of kainic acid-treated fosB null

mice.

Brain Res Mol Brain Res 1997 Jul;47(1-2):31-43

Hollen KM, Nakabeppu Y, Davies SW.

Changes in expression of delta FosB and the Fos family proteins following NMDA receptor activation in the rat

striatum.

Oncogene 1995 Apr 6:10(7):1343-51

Oda S, Nishida J, Nakabeppu Y, Sekiguchi M.

Stabilization of cyclin E and cdk2 mRNAs at G1/S transition in Rat-1A cells emerging from the G0 state.

Neuron 1994 Nov;13(5):1235-44

Hope BT, Nye HE, Kelz MB, Self DW, ladarola MJ, Nakabeppu Y, Duman RS, Nestler EJ. Induction of a long-lasting AP-1 complex composed of altered Fos-like proteins in brain by chronic cocaine and

other chronic treatments.

D4573.C38

The state of the s

From:

Woitach, Joseph

Sent:

Tuesday, May 06, 2003 10:55 AM

To: Subject: STIC-ILL references

Hello.

Could you send me the following references:

Cell 1991 Feb 22;64(4):751-9 Nakabeppu Y, Nathans D.

A naturally occurring truncated form of FosB that inhibits Fos/Jun transcriptional activity.

Mol Cell Biol 1993 Jul;13(7):4157-66

Nakabeppu Y, Oda S, Sekiguchi M.

Proliferative activation of quiescent Rat-1A cells by delta FosB

J Neurochem 1997 Jan;68(1):272-9

Bing G, Wang W, Qi Q, Feng Z, Hudson P, Jin L, Zhang W, Bing R, Hong JS. Long-term expression of Fos-related antigen and transient expression of delta FosB associated with seizures in the

rat hippocampus and striatum.

Eur J Neurosci 1998 May;10(5):1743-51

Porcella A. Gessa GL, Pani L.

Delta9-tetrahydrocannabinol increases sequence-specific AP-1 DNA-binding activity and Fosrelated antigens in

the rat brain.

Brain Res Mol Brain Res 1998 Jan;53(1-2):41-52

Perez-Otano I, Mandelzys A, Morgan JI.

MPTP-Parkinsonism is accompanied by persistent expression of a delta-FosB-like protein in dopaminergic

pathways.

J Neurosci. 1997 Jul 15;17(14):5407-15

Mandelzys A, Gruda MA, Bravo R, Morgan JI.

Absence of a persistently elevated 37 kDa fos-related antigen and AP-1-like DNA-binding activity in the brains of kainic acid-treated fosB null mice.

Brain Res Mol Brain Res 1997 Jul:47(1-2):31-43

Hollen KM, Nakabeppu Y, Davies SW.

Changes in expression of delta FosB and the Fos family proteins following NMDA receptor activation in the rat

striatum.

Oncogene 1995 Apr 6;10(7):1343-51

Oda S, Nishida J, Nakabeppu Y, Sekiguchi M.

Stabilization of cyclin E and cdk2 mRNAs at G1/S transition in Rat-1A cells emerging from the G0 state.

Neuron 1994 Nov;13(5):1235-44

Hope BT, Nye HE, Kelz MB, Self DW, ladarola MJ, Nakabeppu Y, Duman RS, Nestler EJ. Induction of a long-lasting AP-1 complex composed of altered Fos-like proteins in brain by chronic cocaine and

other chronic treatments.

Woitach, Joseph

From:

Woitach, Joseph

Sent:

Tuesday, May 06, 2003 10:55 AM

To: Subject: STIC-ILL references

Hello.

Could you send me the following references:

Cell 1991 Feb 22;64(4):751-9

Nakabeppu Y, Nathans D.

A naturally occurring truncated form of FosB that inhibits Fos/Jun transcriptional activity.

Mol Cell Biol 1993 Jul;13(7):4157-66

Nakabeppu Y, Oda S, Sekiguchi M.

Proliferative activation of quiescent Rat-1A cells by delta FosB

J Neurochem 1997 Jan;68(1):272-9

Bing G, Wang W, Qi Q, Feng Z, Hudson P, Jin L, Zhang W, Bing R, Hong JS.

Long-term expression of Fos-related antigen and transient expression of delta FosB associated with seizures in the

rat hippocampus and striatum.

Eur J Neurosci 1998 May;10(5):1743-51

Porcella A, Gessa GL, Pani L.

Delta9-tetrahydrocannabinol increases sequence-specific AP-1 DNA-binding activity and Fosrelated antigens in

the rat brain.

Brain Res Mol Brain Res 1998 Jan;53(1-2):41-52

Perez-Otano I, Mandelzys A, Morgan JI.

MPTP-Parkinsonism is accompanied by persistent expression of a delta-FosB-like protein in dopaminergic

pathways.

J Neurosci. 1997 Jul 15;17(14):5407-15

Mandelzys A, Gruda MA, Bravo R, Morgan JI.

Absence of a persistently elevated 37 kDa fos-related antigen and AP-1-like DNA-binding activity in the brains of kainic acid-treated fosB null

mice.

Brain Res Mol Brain Res 1997 Jul;47(1-2):31-43

Hollen KM, Nakabeppu Y, Davies SW.

Changes in expression of delta FosB and the Fos family proteins following NMDA receptor activation in the rat

striatum.

Oncogene 1995 Apr 6;10(7):1343-51

Oda S, Nishida J, Nakabeppu Y, Sekiguchi M.

Stabilization of cyclin E and cdk2 mRNAs at G1/S transition in Rat-1A cells emerging from the

G0 state.

Neuron 1994 Nov;13(5):1235-44

Hope BT, Nye HE, Kelz MB, Self DW, ladarola MJ, Nakabeppu Y, Duman RS, Nestler EJ. Induction of a long-lasting AP-1 complex composed of altered Fos-like proteins in brain by chronic cocaine and

other chronic treatments.

Thank you,

Joseph Woitach (703) 305-3732

office CM1/12E05 mail CM1/12E12

Tracking:

Recipient

Delivery

Read

STIC-ILL

Delivered: 5/6/03 10:55 AM

Read: 5/6/03 10:58 AM